§4.113

DISEASES OF THE HEART—Continued

Rating

10

NOTE: These evaluations are for involvement of a single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.

7122 Cold injury residuals:

With the following in affected parts:

Arthralgia or other pain, numbness, or cold sensitivity plus two or more of the following: tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis)

Arthralgia or other pain, numbness, or cold sensitivity plus tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, or X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis)

Arthralgia or other pain, numbness, or cold sensitivity

NOTE (1): Separately evaluate amputations of fingers or toes, and complications such as squamous cell carcinoma at the site of a cold injury scar or peripheral neuropathy, under other diagnostic codes. Separately evaluate other disabilities that have been diagnosed as the residual effects of cold injury, such as Raynaud's phenomenon, muscle atrophy, etc., unless they are used to support an evaluation under diagnostic code 7122.

NOTE (2): Evaluate each affected part (e.g., hand, foot, ear, nose) separately and combine the ratings in accordance with §§ 4.25 and 4.26.

7123 Soft tissue sarcoma (of vascular origin) ...

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.

(Authority: 38 U.S.C. 1155)

 $[62\ FR\ 65219,\ Dec.\ 11,\ 1997,\ as\ amended\ at\ 63\ FR\ 37779,\ July\ 14,\ 1998;\ 71\ FR\ 52460,\ Sept.\ 6,\ 2006;\ 79\ FR\ 2100,\ Jan.\ 13,\ 2014;\ 82\ FR\ 50804,\ Nov.\ 2,\ 2017]$

THE DIGESTIVE SYSTEM

§ 4.110 Ulcers.

Experience has shown that the term "peptic ulcer" is not sufficiently specific for rating purposes. Manifest differences in ulcers of the stomach or duodenum in comparison with those at an

anastomotic stoma are sufficiently recognized as to warrant two separate graduated descriptions. In evaluating the ulcer, care should be taken that the findings adequately identify the particular location.

§4.111 Postgastrectomy syndromes.

There are various postgastrectomy symptoms which may occur following anastomotic operations of the stomach. When present, those occurring during or immediately after eating and known as the "dumping syndrome" are characterized by gastrointestinal complaints and generalized symptoms simulating hypoglycemia; those occurring from 1 to 3 hours after eating usually present definite manifestations of hypoglycemia.

§ 4.112 Weight loss.

For purposes of evaluating conditions in §4.114, the term "substantial weight loss" means a loss of greater than 20 percent of the individual's baseline weight, sustained for three months or longer; and the term "minor weight loss" means a weight loss of 10 to 20 percent of the individual's baseline weight, sustained for three months or longer. The term "inability to gain weight" means that there has been substantial weight loss with inability to regain it despite appropriate therapy. "Baseline weight" means the average weight for the two-year-period preceding onset of the disease.

(Authority: 38 U.S.C. 1155) [66 FR 29488, May 31, 2001]

§4.113 Coexisting abdominal conditions.

There are diseases of the digestive system, particularly within the abdomen, which, while differing in the site of pathology, produce a common disability picture characterized in the main by varying degrees of abdominal distress or pain, anemia and disturbances in nutrition. Consequently, certain coexisting diseases in this area, as indicated in the instruction under the

§4.114

title "Diseases of the Digestive System," do not lend themselves to distinct and separate disability evaluations without violating the fundamental principle relating pyramiding as outlined in §4.14.

§4.114 Schedule of ratings—digestive

Ratings under diagnostic codes 7301 to 7329, inclusive, 7331, 7342, and 7345 to 7348 inclusive will not be combined with each other. A single evaluation will be assigned under the diagnostic code which reflects the predominant disability picture, with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation.

	Rating
7200 Mouth, injuries of.	
Rate as for disfigurement and impairment of	
function of mastication.	
7201 Lips, injuries of.	
Rate as for disfigurement of face.	
7202 Tongue, loss of whole or part:	
With inability to communicate by speech	100
One-half or more	60
With marked speech impairment	30
7203 Esophagus, stricture of:	
Permitting passage of liquids only, with marked	
impairment of general health	80
Severe, permitting liquids only	50
Moderate	30
7204 Esophagus, spasm of (cardiospasm).	
If not amenable to dilation, rate as for the de-	
gree of obstruction (stricture).	
7205 Esophagus, diverticulum of, acquired.	
Rate as for obstruction (stricture).	
7301 Peritoneum, adhesions of:	
Severe; definite partial obstruction shown by X-	
ray, with frequent and prolonged episodes of	
severe colic distension, nausea or vomiting,	
following severe peritonitis, ruptured appen- dix, perforated ulcer, or operation with drain-	
age	50
Moderately severe; partial obstruction mani-	30
fested by delayed motility of barium meal and	
less frequent and less prolonged episodes of	
pain	30
Moderate; pulling pain on attempting work or	30
aggravated by movements of the body, or	
occasional episodes of colic pain, nausea,	
constipation (perhaps alternating with diar-	
rhea) or abdominal distension	10
Mild	0
Note: Ratings for adhesions will be considered	"
when there is history of operative or other	
traumatic or infectious (intraabdominal) proc-	
ess, and at least two of the following: disturb-	
ance of motility, actual partial obstruction, re-	
flex disturbances, presence of pain.	
7304 Ulcer, gastric.	
7305 Ulcer, duodenal:	
Severe; pain only partially relieved by standard	
ulcer therapy, periodic vomiting, recurrent	
hematemesis or melena, with manifestations	
	I

of anemia and weight loss productive of defi-

nite impairment of health ...

•	
	Rating
Moderately severe; less than severe but with impairment of health manifested by anemia and weight loss; or recurrent incapacitating	
episodes averaging 10 days or more in dura- tion at least four or more times a year	40
days in duration; or with continuous mod- erate manifestations	20
yearly	10
Pronounced; periodic or continuous pain unrelieved by standard ulcer therapy with periodic vomiting, recurring melena or hematemesis, and weight loss. Totally inca-	
pacitating	100
nounced and less continuous symptoms with definite impairment of health	60
dominal pain at least once a month partially or completely relieved by ulcer therapy, mild and transient episodes of vomiting or melena	40
Moderate; with episodes of recurring symptoms several times a year	20
Mild; with brief episodes of recurring symptoms once or twice yearly	10
7307 Gastritis, hypertrophic (identified by gastroscope): Chronic; with severe hemorrhages, or large ul-	
cerated or eroded areas Chronic; with multiple small eroded or ulcerated	60
areas, and symptoms	30
toms	10
ing pernicious anemia. Rate the underlying condition.	
7308 Postgastrectomy syndromes: Severe; associated with nausea, sweating, cir- culatory disturbance after meals, diarrhea, hypoglycemic symptoms, and weight loss	
with malnutrition and anemia	60
weight loss	40
with characteristic mild circulatory symptoms or continuous mild manifestations	20
Rate as for gastric ulcer. 7310 Stomach, injury of, residuals.	
Rate as peritoneal adhesions. 7311 Residuals of injury of the liver:	
Depending on the specific residuals, separately evaluate as adhesions of peritoneum (diagnostic code 7301), cirrhosis of liver (diagnostic code 7312), and chronic liver disease without cirrhosis (diagnostic code 7345).	
7312 Cirrhosis of the liver, primary biliary cirrhosis, or cirrhotic phase of sclerosing cholangitis:	
Generalized weakness, substantial weight loss, and persistent jaundice, or; with one of the following refractory to treatment: ascites, he- patic encephalopathy, hemorrhage from	
varices or portal gastropathy (erosive gastritis)	100
•	

60